

Cross-resistance and biochemical mechanisms of resistance to indoxacarb in the diamondback moth, *Plutella xylostella*



Shuzhen Zhang¹, Xiaolei Zhang¹, Jun Shen, Dongyang Li, Hu Wan, Hong You^{*}, Jianhong Li^{*}

Hubei Insect Resources Utilization and Sustainable Pest Management Key Laboratory, College of Plant Science and Technology, Huazhong Agricultura

Studies of cross-resistance, resistance stability and metabolic detoxification mechanisms of laboratory-selected indoxacarb resistant strains are important strategies to delay indoxacarb resistance evolution in field populations of *P. xylostella*. In the present study, cross-resistance to metaflumizone, beta-cypermethrin, chlorfenapyr and other insecticides was determined in a laboratory-selected resistant strain of *P. xylostella* with a high level of indoxacarb resistance. Resistance stability was also studied in the resistant strain. Furthermore, the effects of three synergists on indoxacarb toxicity were tested to confirm the involvement of metabolic mechanisms in indoxacarb resistance in *P. xylostella*. Finally, metabolic enzymes were investigated in the resistant strain. The study provided important information on indoxacarb resistance characteristics that will be useful for the indoxacarb resistance management of *P. xylostella*.

2. Materials and methods

2.1. Susceptible and resistant strains

The susceptible strain of *P. xylostella*, obtained from Wuhan in 2002, has been maintained in the laboratory for >10 years without exposure to any insecticides. The resistant strain derived from the susceptible strain was selected with 1 to 500 mg/L of indoxacarb for 61 generations. The mortality during selection cycles was approximately 30–70%. Adults were fed with 10% honey solution and allowed to lay eggs on radish seedlings (*Raphanus sativus* L.). All populations were maintained at 25 ± 1 °C and 50–70% relative humidity with a 16:8 h light:dark photoperiod.

2.2. Insecticides

Six commonly used insecticides, including abamectin (2% Emulsifiable Concentrate (EC)), chlorfluazuron (5% EC), chlorfenapyr (10% EC), diafenthiuron (20% EC), beta-cypermethrin (20% EC) and chlorantraniliprole (5% EC), were provided by the Institute of Plant Protection of the Guangdong Academy of Agricultural Sciences, China. Cyantraniliprole (10% Suspension Concentrate (SC)) was supplied by E.I. DuPont Co. Spinosad (25 g/L SC) was supplied by the Dow AgroSciences Company. The indoxacarb (96%) and metaflumizone (97%) insecticides used in this study were technical grade compounds and were supplied by the Hubei Kangbaotai Fine-Chemicals Co., Ltd. Reagent grade triphenyl phosphate (TPP), diethyl maleate (DEM) and piperonyl butoxide (PBO) were purchased from Sigma.

2.3. Bioassay

Insecticide toxicity was assayed using the leaf-dipping method as described previously with a slight modification [21]. Young cabbage (*Brassica oleracea*) leaf discs (6.5 cm diameter) were cut from plants grown in a greenhouse without any insecticides. The technical grade compounds of indoxacarb and metaflumizone were dissolved in *N,N*-dimethylformamide, and then were serially diluted to five to seven required concentrations with distilled water containing 0.1% Triton X-100. Other insecticides were serially diluted to five to seven required concentrations with distilled water containing 0.1% Triton X-100. Three leaf discs were grouped together and dipped in solutions with different insecticide concentrations for 10 s. Control discs were only treated with 0.1% Triton X-100 solution and the control mortality was required to be <10%. All dipped leaf discs were dried at room temperature for approximately 120 min. The discs were then placed individually in plastic petri dishes (7.0 cm diameter). A total of 10 third-instar larvae were transferred to each dish, and four replicates were prepared and kept under standard conditions as previously described [10]. For the analysis of synergistic effects, 100 mg/L of DEM, TPP, and PBO were prepared. Larvae mortality was recorded after 48 h for abamectin,

spinosad, chlorfenapyr, diafenthiuron, indoxacarb, and beta-cypermethrin, and after 72 h for chlorfluazuron, cyantraniliprole, and chlorantraniliprole. Larvae were considered dead if they could not be induced to move when touched with a brush.

2.4. Enzyme activity

Esterase activity was determined by the method of Asperen with modification [22]. The enzyme source was the supernatant of 10 third-instar larvae homogenized on ice in 1000 μ L of precooled homogenization buffer (0.04 M sodium phosphate buffer, pH 7.0), then centrifuged at 4 °C and $10,000 \times g$ for 15 min. To determine esterase activity, 1000 μ L of preheated 0.3 mM p-NA and 200 μ L of enzyme source (diluted 20–100-fold) were added to Eppendorf tubes; after 15 min, the dyeing reagent was added, and after 30 min, the optical density (OD) at 600 nm was recorded with an ultraviolet spectrophotometer.

GST activity using CDNB as substrates was measured as follows [23]. The enzyme source was prepared with 30 third-instar larvae homogenized in 1 mL of homogenization buffer (0.1 M sodium phosphate buffer with 1 mM EDTA, pH 6.5). The reaction solutions contained 50 μ L of enzyme, 790 μ L of the homogenization buffer, 30 μ L of 30 mM CDNB and 30 μ L of 30 mM GSH. An enzyme solution instead by buffer served as control. The OD at 340 nm was recorded at intervals of 1 s for 2 min at 25 °C with an ultraviolet spectrophotometer.

Cytochrome P450 monooxygenase activity was determined as follows [24]. The enzyme source was obtained from the homogenization of 50 third-instar larvae on ice in 1 mL of 0.1 M sodium phosphate buffer (pH 7.5) containing 1 mM of EDTA, 1 mM of DTT, 1 mM of PTU and 1 mM of PMSF, and centrifuged at $18,000 \times g$ for 30 min. Afterwards, 600 μ L of supernatant was added to an Eppendorf tube. The supernatant was recentrifuged at $18,000 \times g$ for 30 min. The supernatant was then used as the substrate to test cytochrome P450 monooxygenase activity. Quantities of 750 μ L of 2 mM PNA, 75 μ L of 9.6 mM NADPH and 675 μ L of enzyme solution were added to Eppendorf tubes. The mixed solutions were warmed in a water bath at 34 °C for 30 min, and the OD at 405 nm was recorded with an ultraviolet spectrophotometer.

The protein content of the enzyme solutions was determined by the Bradford method: 100 μ L of sample, 700 μ L of sterilized deionized water and 200 μ L of protein assay dye reagent were added to Eppendorf tubes; after 5 min, the OD at 595 nm was recorded with an ultraviolet spectrophotometer.

2.5. Indoxacarb resistance stability in *P. xylostella*

Third-instar larvae from the resistant strain that had been maintained for seven generations in the absence of selection pressure, were bioassayed at $F_{53'}$, $F_{55'}$, $F_{57'}$ and $F_{58'}$. The resistance ratio (RR) for the unselected populations was estimated by dividing their LC_{50} values by the LC_{50} value of the susceptible strain.

2.6. Statistical analysis

The mortality data were corrected by the control mortality using Abbott's formula; LC_{50} values, 95% confidence intervals and slopes were calculated by Probit analysis. The resistance ratio (RR) was calculated by dividing the LC_{50} value of the resistant strain by the LC_{50} value of the susceptible strain (Tables 1 and 4). The classification of the resistance levels followed Shao et al. [21]: $RR \leq 5$ -fold was classified as susceptible, $RR = 5$ –10-fold as a low level of resistance, $RR = 10$ –100-fold as a medium level of resistance and $RR > 100$ -fold as a high level of resistance.

Table 1
Indoxacarb resistance changes in the laboratory population of *P. xylostella* with continuing selection.

| Generation | No. | LC ₅₀ (95% CL) mg/L | Slope(SE) | ² (df) | RR |
|-----------------|-----|--------------------------------|-------------|-------------------|--------|
| Sus. | 240 | 0.55 (0.35–0.83) | 1.26 (0.27) | 0.40 (3) | 1.0 |
| F ₃₁ | 240 | 9.45 (6.51–17.48) | 1.51 (0.31) | 1.72 (3) | 17.2 |
| F ₃₂ | 240 | 9.63 (4.97–65.32) | 0.93 (0.27) | 0.02 (2) | 17.5 |
| F ₃₃ | 240 | 3.02 (0.91–5.34) | 1.08 (0.29) | 0.47 (2) | 5.5 |
| F ₃₆ | 240 | 3.93 (2.36–5.86) | 1.40 (0.30) | 0.52 (3) | 7.2 |
| F ₃₈ | 240 | 15.56 (10.18–32.22) | 1.47 (0.27) | 0.95 (2) | 28.3 |
| F ₄₃ | 240 | 3.35 (2.27–5.18) | 1.31 (0.27) | 1.78 (2) | 6.1 |
| F ₄₄ | 240 | 23.26 (12.36–90.76) | 0.96 (0.22) | 0.38 (3) | 42.3 |
| F ₄₅ | 240 | 44.84 (23.30–205.07) | 0.93 (0.23) | 0.49 (3) | 81.5 |
| F ₄₇ | 240 | 17.08 (12.17–24.95) | 1.33 (0.21) | 2.21 (2) | 31.1 |
| F ₄₈ | 240 | 25.69 (16.11–52.67) | 1.06 (0.22) | 0.59 (2) | 46.7 |
| F ₅₀ | 240 | 14.36 (9.55–20.56) | 1.33 (0.22) | 1.83 (3) | 26.1 |
| F ₅₁ | 240 | 8.71 (4.67–13.39) | 1.24 (0.21) | 3.39 (3) | 15.8 |
| F ₅₃ | 240 | 46.95 (35.07–62.42) | 1.63 (0.19) | 1.45 (3) | 85.4 |
| F ₅₄ | 240 | 68.72 (47.09–104.97) | 1.84 (0.23) | 0.41 (3) | 124.9 |
| F ₅₅ | 240 | 63.46 (44.86–88.01) | 1.44 (0.21) | 0.89 (2) | 115.4 |
| F ₅₇ | 240 | 145.43 (105.65–219.74) | 1.50 (0.22) | 4.18 (2) | 264.4 |
| F ₅₈ | 240 | 588.13 (312.64–2440.98) | 1.00 (0.24) | 0.85 (3) | 1069.3 |
| F ₆₀ | 240 | 538.55 (298.76–1727.76) | 0.87 (0.24) | 0.94 (2) | 979.2 |
| F ₆₁ | 240 | 574.12 (389.54–932.45) | 1.19 (0.20) | 2.21 (3) | 1043.9 |

3. Results

3.1. Resistance development assessment

The resistant strain of *P. xylostella* (IR) was derived in the laboratory from a susceptible strain (IS) following 61 generations of continuous indoxacarb selection (Table 1). After 53 generations, the IR had evolved only an 85.4-fold resistance to indoxacarb. Afterwards, the RR increased quickly from the 54th to the 58th generation, reaching a 1069.3-fold increase by the 58th generation, and showed no further changes from the 58th to the 61st generation. Additionally, indoxacarb resistance changes in *P. xylostella* with cessation of selection were monitored when the RR was 85.4-fold. After cessation of selection, the RR showed a quick 12.5-fold decline from F₅₃' to F₅₈' (Table 2).

3.2. Cross-resistance

Compared with the IS, the IR showed cross-resistance to three insecticides, including beta-cypermethrin (RR = 8.7), metaflumizone (RR = 20.9), and chlorfenapyr (RR = 8.3). No significant cross-resistance to other insecticides was observed (Table 3).

3.3. Synergistic effects

The synergistic effects of PBO, TPP, and DEM with indoxacarb against the IR and the IS are shown in Table 4. Significant synergisms were found for PBO and DEM in the IR (F₅₈) (synergistic ratio, SR = 7.8- and 3.5-fold, respectively); no synergy was observed for TPP in either strain.

Table 2
Indoxacarb resistance changes in the laboratory population of *P. xylostella* with cessation of selection.

| Generation | No. | LC ₅₀ (95% CL) mg/L | Slope (SE) | ² (df) | RR |
|-------------------|-----|--------------------------------|-------------|-------------------|------|
| F ₅₃ ' | 240 | 46.95 (35.07–62.42) | 1.63 (0.19) | 1.45 (3) | 85.4 |
| F ₅₅ ' | 240 | 31.72 (22.41–45.09) | 1.63 (0.29) | 0.80 (2) | 57.7 |
| F ₅₇ ' | 240 | 13.33 (9.61–18.08) | 1.87 (0.37) | 0.28 (2) | 24.2 |
| F ₅₈ ' | 240 | 6.85 (5.45–9.07) | 2.42 (0.34) | 3.55 (3) | 12.5 |

Table 3
Cross-resistance spectrum in laboratory indoxacarb-resistant strains of *P. xylostella*.

| Insecticides | Strains | LC ₅₀ (95% CL) mg/L | Slope (SE) | ² (df) | RR |
|---------------------|-----------------|--------------------------------|-------------|-------------------|------|
| Beta-cypermethrin | S | 5.41 (3.80–19.67) | 1.88 (0.58) | 3.13 (2) | |
| | F ₅₃ | 47.28 (28.54–97.17) | 1.08 (0.26) | 0.51 (3) | 8.7 |
| Metaflumizone | S | 1.24 (0.92–1.62) | 2.16 (0.40) | 1.51 (3) | |
| | F ₅₃ | 25.90 (14.57–52.48) | 1.12 (0.26) | 0.89 (3) | 20.9 |
| Diafenthiuron | S | 20.83 (15.28–27.19) | 1.85 (0.24) | 1.60 (2) | |
| | F ₅₇ | 25.47 (17.12–34.13) | 1.84 (0.30) | 3.16 (2) | 1.2 |
| Chlorfenapyr | S | 0.36 (0.24–0.57) | 1.38 (0.29) | 2.70 (3) | |
| | F ₅₈ | 2.99 (2.51–3.71) | 3.87 (0.59) | 1.26 (3) | 8.3 |
| Spinosad | S | 0.54 (0.36–1.00) | 1.50 (0.27) | 2.78 (3) | |
| | F ₅₈ | 0.86 (0.62–1.52) | 2.07 (0.39) | 2.00 (3) | 1.6 |
| Chlorantraniliprole | S | 0.22 (0.16–0.31) | 1.68 (0.23) | 1.23 (2) | |
| | F ₅₈ | 0.21 (0.14–0.34) | 1.47 (0.30) | 0.08 (3) | 1.0 |
| Chlorfluazuron | S | 1.04 (0.45–370.12) | 1.21 (0.49) | 0.08 (3) | |
| | F ₅₈ | 1.00 (0.70–1.83) | 1.63 (0.32) | 3.53 (2) | 1.0 |
| Abamectin | S | 0.05 (0.04–0.07) | 1.45 (0.21) | 0.41 (3) | |
| | F ₅₈ | 0.04 (0.03–0.08) | 2.29 (0.43) | 0.49 (2) | 0.8 |
| Cyantraniliprole | S | 0.20 (0.15–0.28) | 2.39 (0.29) | 0.72 (3) | |
| | F ₅₈ | 0.19 (0.14–0.26) | 1.98 (0.32) | 3.63 (3) | 1.0 |

3.4. Metabolic enzyme activity

Esterase activity of the IR (F₅₈) was 1.47 mol/min/mg protein, with a 1.6-fold increase over that of the IS (0.93 mol/min/mg protein) (Table 5). Cytochrome P450 monooxygenase activity using p-nitroanisole as substrate in the IS and IR is shown in Table 5. The cytochrome P450 monooxygenase activity of the IR (F₅₈) was 6.8-fold compared with that of the IS. GST activity of the IR (F₅₈) was 5.8-fold as that of the IS.

4. Discussion

Indoxacarb belongs to a relatively new class of sodium channel blocker insecticides [17]. Due to the intensive use of indoxacarb, it had been reported that resistance had occurred in field populations of some insect species, such as *Aedes albopictus*, *Choristoneura rosaceana*, *Earias vittella*, *Helicoverpa armigera*, *Heliothis virescens*, *Lobesia botrana*, *Musca domestica*, *Sitophilus zeamais*, *Spodoptera exigua*, *Kampimodromus aberrant*, *Spodoptera litura*, *Tuta absoluta* and *P. xylostella* [10,13,17,18,25–35]. Currently, a high level of resistance to indoxacarb in *P. xylostella* is common in South China [17], whereas only low and moderate levels of resistance are found in Central China [10]. Thus, to delay indoxacarb resistance evolution in field populations of *P. xylostella*, studies concerning resistance stability, cross-resistance pattern as well as resistance mechanisms in this species are very important to develop sustainable pest management strategies in the field.

Previous studies have shown that *P. xylostella* can maintain resistance to indoxacarb under selection pressure in the laboratory [7]. Continuous selection resulted in a high level of resistance from an 813-fold resistance in the 1st generation to a 2594-fold resistance by the 8th generation [7]. This resistance was unstable and disappeared after a few generations without insecticide pressure [7]. Similarly, in the present study, *P. xylostella* selected with indoxacarb for 58 generations in the laboratory developed a 1069.3-fold resistance. The resistance was also unstable and disappeared after a few generations without insecticide pressure. This instability might be associated with an incompletely recessive trait of indoxacarb resistance and fitness costs [36]. An unstable resistance is useful to control insecticide resistance by strategically stopping the use of an insecticide to prolong its efficiency.

Cross-resistance between indoxacarb and other insecticides has been reported in *P. xylostella* strains in several studies [1,37]. Some reports indicated that cross-resistance could result from metabolic detoxification enzymes (cytochrome P450 monooxygenase, GST and esterase), mutation at an insecticidal target site or delayed cuticular penetration [38]. In the present study, there was cross-resistance between indoxacarb and metaflumizone in the IR. Metaflumizone belongs

Table 4
Synergism of TPP, DEM and PBO on indoxacarb in different strains of *P. xylostella*.

| Generation | Insecticides | LC ₅₀ (95% CL) mg/L | Slope (SE) | ² (df) | SR |
|-----------------|------------------|--------------------------------|-------------|-------------------|-----|
| Unselected | Indoxacarb | 0.72 (0.51–0.95) | 1.87 (0.30) | 0.17 (3) | 1.0 |
| | Indoxacarb + PBO | 1.50 (1.05–2.50) | 1.52 (0.24) | 0.31 (2) | 0.5 |
| | Indoxacarb + DEM | 0.92 (0.70–1.20) | 12.7 (0.45) | 0.33 (2) | 0.8 |
| | Indoxacarb + TPP | 0.72 (0.56–1.00) | 2.09 (0.32) | 0.68 (3) | 1.0 |
| F ₄₇ | Indoxacarb | 17.08 (12.17–24.95) | 1.33 (0.21) | 2.21 (2) | 1.0 |
| | Indoxacarb + PBO | 15.32 (11.14–23.89) | 1.58 (0.29) | 0.99 (2) | 1.1 |
| | Indoxacarb + DEM | 18.16 (12.38–28.76) | 1.53 (0.25) | 0.17 (2) | 0.9 |
| | Indoxacarb + TPP | 11.60 (7.51–18.12) | 1.56 (0.23) | 1.86 (3) | 1.5 |
| F ₅₈ | Indoxacarb | 588.13 (312.64–2440.98) | 1.00 (0.24) | 0.85 (3) | 1.0 |
| | Indoxacarb + PBO | 75.73 (45.62–129.11) | 1.17 (0.24) | 1.49 (3) | 7.8 |
| | Indoxacarb + DEM | 170.68 (125.90–268.40) | 1.92 (0.34) | 1.33 (3) | 3.5 |
| | Indoxacarb + TPP | 400.16 (225.83–2049.03) | 1.80 (0.48) | 0.41 (3) | 1.5 |

to a new chemical class of insecticides with the same mode of action as indoxacarb, a sodium channel blocker in the central nervous system [39–41]. A more recent study also showed that field populations of *P. xylostella* with a high level of resistance to indoxacarb also showed resistance to metaflumizone and that sodium channel mutations may result in resistance against the two insecticides in this species [17]. These results suggest that cross-resistance between the two insecticides in *P. xylostella* can be caused by sodium channel mutations. Furthermore, beta-cypermethrin is a pyrethroid insecticide and chlorfenapyr is a pyrrole insecticide. These two insecticides bind at different target sites, implying that metabolic detoxification might be responsible for cross-resistance between them and indoxacarb.

P. xylostella and other insect pests have evolved mechanisms of insecticide resistance to detoxify or reduce their sensitivity to insecticides [1,5,12,42,43]. Increased detoxification can occur by gene duplication of carboxylesterase and increased transcription of cytochrome P450 monooxygenase and GST [42,44]. Moreover, point mutations in the target site can also reduce insecticide binding [5,42]. However, in many cases, up-regulation of the detoxifying enzymes is the most common mechanism of insecticide resistance [42]. Many studies have shown that esterase, GST and cytochrome P450 monooxygenase play an important role in the detoxifying strategies of insect pests [42,44–48]. Currently, indoxacarb resistance through metabolic detoxification has been reported in *P. xylostella* and other insect pests [7,18,19,49]. For example, a synergism study suggested that indoxacarb resistance in *P. xylostella* was esterase mediated [7]. In another study, GST and esterase were involved in resistance to indoxacarb [49]. It was also reported that indoxacarb resistance in *P. xylostella* was associated with cytochrome P450 monooxygenase but not with esterase or GST [18]. However, in the present study, enhanced activity of GST and cytochrome P450 monooxygenase is likely the main detoxification mechanism responsible for resistance by biochemical analysis and synergistic suppression (PBO and DEM).

This study demonstrates that *P. xylostella* can develop resistance to indoxacarb and that enhanced detoxification by GST and cytochrome P450 monooxygenase is mainly responsible for this resistance. The study represents a first step towards understanding the biochemical mechanisms of indoxacarb resistance in selected strains in our laboratory. The target insensitivity mechanism accompanying the appearance of a higher resistance level will be the focus of further investigation. The inheritance patterns of indoxacarb resistance in *P. xylostella* will also

require additional research to confirm the resistance mechanism and to establish more efficient management strategies for this pest.

Competing interests

The authors have declared that no competing interests exist.

Acknowledgments

This work was supported by grants from the National Key Research and Development Program of China (2016YFD0200500 and 2016YED0201008).

References

- [1] M.J. Furlong, D.J. Wright, L.M. Dossall, Diamondback moth ecology and management: problems, progress, and prospects, *Annu. Rev. Entomol.* 58 (2013) 517–541.
- [2] Z.Y. Li, X. Feng, S.S. Liu, M.S. You, M.J. Furlong, Biology, ecology, and management of the Diamondback Moth in China, *Annu. Rev. Entomol.* 61 (2016) 277–296.
- [3] M.P. Zalucki, A. Shabbir, R. Silva, D. Adamson, S.S. Liu, M.J. Furlong, Estimating the economic cost of one of the world's major insect pests, *Plutella xylostella* (Lepidoptera: Plutellidae): just how long is a piece of string? *J. Econ. Entomol.* 105 (2012) 1115–1129.
- [4] N.S. Talekar, Biology, ecology, and management of the diamondback moth, *Annu. Rev. Entomol.* 38 (1993) 275–301.
- [5] L. Guo, L. Liang, X.G. Zhou, X.W. Gao, Novel mutations and mutation combinations of ryanodine receptor in a chlorantraniliprole resistant population of *Plutella xylostella* (L.), *Sci. Rep-UK* 4 (2014) 6924.
- [6] X.L. Wang, S.K. Khakame, C. Ye, Y.H. Yang, Y.D. Wu, Characterisation of field-evolved resistance to chlorantraniliprole in the diamondback moth, *Plutella xylostella*, from China, *Pest Manag. Sci.* 69 (2013) 661–665.
- [7] A.H. Sayyed, D.J. Wright, Genetics and evidence for an esterase-associated mechanism of resistance to indoxacarb in a field population of diamondback moth (Lepidoptera: Plutellidae), *Pest Manag. Sci.* 62 (2006) 1045–1051.
- [8] J.Z. Zhao, Y.X. Li, H.L. Collins, L. Gusukuma-Minuto, R.F.L. Mau, G.D. Thompson, A.M. Shelton, Monitoring and characterization of diamondback moth (Lepidoptera: Plutellidae) resistance to spinosad, *J. Econ. Entomol.* 95 (2002) 430–436.
- [9] L. Zhou, J. Huang, H. Xu, Monitoring resistance of field populations of diamondback moth *Plutella xylostella* L. (Lepidoptera: Yponomeutidae) to five insecticides in South China: a ten-year case study, *Crop. Prot.* 30 (2011) 272–278.
- [10] S.Z. Zhang, X.L. Zhang, J. Shen, K.K. Mao, H. You, J.H. Li, Susceptibility of field populations of the diamondback moth, *Plutella xylostella*, to a selection of insecticides in Central China, *Pestic. Biochem. Physiol.* 132 (2016) 38–46.
- [11] Y.M. Xia, Y.H. Lu, J. Shen, X.W. Gao, H. Qiu, J.H. Li, Resistance monitoring for eight insecticides in *Plutella xylostella* in central China, *Crop. Prot.* 63 (2014) 131–137.
- [12] T.T. Jiang, S.F. Wu, T.T. Yang, C. Zhu, C.F. Gao, Monitoring field populations of *Plutella xylostella* (Lepidoptera: Plutellidae) for resistance to eight insecticides in China, *Fla. Entomol.* 98 (2015) 65–73.
- [13] APRD, Arthropod Pesticide Resistance Database, www.pesticideresistance.com 2016 [Date of access: 02/04/2016].
- [14] J.Z. Zhao, H.L. Collins, Y.X. Li, R.F.L. Mau, G.D. Thompson, M. Hertlein, J.T. Andaloro, R. Boykin, A.M. Shelton, Monitoring of diamondback moth (Lepidoptera: Plutellidae) resistance to spinosad, indoxacarb, and emamectin benzoate, *J. Econ. Entomol.* 99 (2006) 176–181.
- [15] S.F. McCann, G.D. Annis, R. Shapiro, D.W. Piotrowski, G.P. Lahm, J.K. Long, K.C. Lee, M.M. Hughes, B.J. Myers, S.M. Griswold, B.M. Reeves, R.W. March, P.L. Sharpe, P. Lowder, W.E. Barnette, K.D. Wing, The discovery of indoxacarb: oxadiazines as a new class of pyrazoline-type insecticides, *Pest Manag. Sci.* 57 (2001) 153–164.
- [16] B. Lapiet, F.O. Grolleau, D.B. Sattelle, Indoxacarb, an oxadiazine insecticide, blocks insect neuronal sodium channels, *Brit. J. Pharmacol.* 132 (2001) 587–595.

Table 5
Mean (\pm standard error) metabolic enzyme activity in populations of *P. xylostella*.

| Strains | Esterase mol/mg pro/min | GST mol/mg pro/min | Cytochrome P450 monooxygenase nmol/mg pro/min |
|-----------------|-------------------------------|--------------------------|---|
| Sus. | 0.93 \pm 0.03b | 0.73 \pm 0.01c | 0.16 \pm 0.002b |
| F ₄₇ | 0.76 \pm 0.03b | 0.86 \pm 0.01b | 0.20 \pm 0.005b |
| F ₅₈ | 1.47 \pm 0.54a | 4.25 \pm 0.24a | 1.09 \pm 0.014a |

- [17] X.L. Wang, W. Su, J.H. Zhang, Y.H. Yang, K. Dong, Y.D. Wu, Two novel sodium channel mutations associated with resistance to indoxacarb and metaflumizone in the diamondback moth, *Plutella xylostella*, *Insect Sci.* 23 (2016) 50–58.
- [18] T. Shono, L. Zhang, J.G. Scott, Indoxacarb resistance in the house fly, *Musca domestica*, *Pestic. Biochem. Physiol.* 80 (2004) 106–112.
- [19] M. Ahmad, R.M. Hollingworth, Synergism of insecticides proves evidence of metabolic mechanisms of resistance in the obliquebanded leafroller *Choristoneura rosaceana* (Lepidoptera: Tortricidae), *Pest Manag. Sci.* 65 (2004) 465–473.
- [20] L.J. Bird, Genetics, cross-resistance and synergism of indoxacarb resistance in *Helicoverpa armigera* (Lepidoptera: Noctuidae), *Pest Manag. Sci.* 73 (2017) 575–581.
- [21] Z.R. Shao, X. Feng, S. Zhang, Z.Y. Li, J.D. Huang, H. Chen, Z.D. Hu, Guideline for Insecticide Resistance Monitoring of *Plutella xylostella* (L.) on Cruciferous Vegetables, China Agriculture Press, Beijing, 2013.
- [22] K.V. Asperen, A study of housefly esterases by means of a sensitive colorimetric method, *J. Insect Physiol.* 8 (1962) 401–414.
- [23] L. Xu, M. Wu, Z.J. Han, Biochemical and molecular characterization and cross-resistance in field and laboratory chlorpyrifos-resistant strains of *Laodelphax striatellus* (Hemiptera: Delphacidae) from eastern China, *Pest Manag. Sci.* 70 (2014) 1118–1129.
- [24] Y.C. Han, W.T. Yu, W.Q. Zhang, Y.H. Yang, T. Walsh, J.G. Oakeshott, Y.D. Wu, Variation in P450-mediated fenvalerate resistance levels is not correlated with CYP37B3 genotype in Chinese populations of *Helicoverpa armigera*, *Pestic. Biochem. Physiol.* 121 (2015) 129–135.
- [25] M. Smirle, T. Lowery, C. Zuruwski, Resistance and cross-resistance to four insecticides in populations of obliquebanded leafroller (Lepidoptera: Tortricidae), *J. Econ. Entomol.* 95 (2002) 820–825.
- [26] A.H. Sayyed, M. Ahmad, N. Crickmore, Fitness cost limit the development of resistance to indoxacarb and deltamethrin in *Heliothis virescens* (Lepidoptera: Noctuidae), *J. Econ. Entomol.* 101 (2008) 1927–1933.
- [27] M. Ahmad, M.I. Arif, Resistance of Pakistani field populations of spotted bollworm *Earias vitella* (Lepidoptera: Noctuidae) to pyrethroid, organophosphorus and new chemical insecticides, *Pest Manag. Sci.* 65 (2009) 433–439.
- [28] G. Silva, M. Picano, L. Bacci, A. Crespo, J. Rosado, R. Guedes, Control failure likelihood and spatial dependence of insecticide resistance in the tomato pinworm, *Tuta absoluta*, *Pest Manag. Sci.* 67 (2011) 913–920.
- [29] H. Khan, W. Akram, K. Shehzad, E. Shaalan, First report of field evolved resistance to agrochemicals in dengue mosquito, *Aedes albopictus* (Diptera: Culicidae), from Pakistan, *Parasite Vectors* 4 (2011) 146.
- [30] H. Tong, Q. Su, X. Zhou, L. Bai, Field resistance of *Spodoptera litura* (Lepidoptera: Noctuidae) to organophosphates, pyrethroids, carbamates and four newer chemistry insecticides in Hunan, China, *J. Pest. Sci.* 86 (2013) 599–609.
- [31] M. Gao, X. Li, W. Mu, P. Zhang, C. Zhou, Resistant levels of *Spodoptera exigua* to eight various insecticides in Shandong, China, *J. Pestic. Sci.* 39 (2014) 7–13.
- [32] S. Civolani, M. Boselli, A. Butturini, M. Chicca, E.A. Fano, S. Cassanelli, Assessment of insecticide resistance of *Lobesia botrana* (Lepidoptera: Tortricidae) in Emilia-Romagna region, *J. Econ. Entomol.* 107 (2014) 1245–1249.
- [33] N. Abbas, S. Shad, M. Ismail, Resistance to conventional and new insecticides in house flies (Diptera: Muscidae) from poultry facilities in Punjab, Pakistan, *J. Econ. Entomol.* 108 (2015) 1–8.
- [34] K. Haddi, L. Mendonca, M.F.D. Santos, R. Guedes, E. Oliveira, Metabolic and behavioral mechanisms of indoxacarb resistance in *Sitophilus zeamais* (Coleoptera: Curculionidae), *J. Econ. Entomol.* 108 (2015) 362–369.
- [35] M. Qayyum, W. Wakil, M. Arif, S. Sahi, N. Saeed, D. Russell, Multiple resistances against formulated organophosphates, pyrethroids, and newer-chemistry insecticides in populations of *Helicoverpa armigera* (Lepidoptera: Noctuidae) from Pakistan, *J. Econ. Entomol.* 108 (2015) 1–8.
- [36] M.B.S. Afzal, S.A. Shad, E. Basoalto, M. Ejaz, J.E. Serrao, Characterization of indoxacarb resistance in *Phenacoccus solenopsis* Tinsley (Homoptera: Pseudococcidae): cross-resistance, stability and fitness cost, *J. Asia Pac. Entomol.* 18 (2015) 779–785.
- [37] A.H. Sayyed, D.J. Wright, Fipronil resistance in the diamondback moth (Lepidoptera: Plutellidae): inheritance and number of genes involved, *J. Econ. Entomol.* 97 (2004) 2043–2050.
- [38] L. Qian, G.C. Cao, J.X. Song, Q. Yin, Z.J. Han, Biochemical mechanisms conferring cross-resistance between tebufenozide and abamectin in *Plutella xylostella*, *Pestic. Biochem. Physiol.* 91 (2008) 175–179.
- [39] V.L. Salgado, J.H. Hayashi, Metaflumizone is a novel sodium channel blocker insecticide, *Vet. Parasitol.* 150 (2007) 182–189.
- [40] K. Takagi, H. Hamaguchi, T. Nishimatsu, T. Konno, Discovery of metaflumizone, a novel semicarbazone insecticide, *Vet. Parasitol.* 150 (2007) 177–181.
- [41] R.T.V. Stein, K.S. Silver, D.M. Soderlund, Indoxacarb, metaflumizone, and other sodium channel inhibitor insecticides: mechanism and site of action on mammalian voltage-gated sodium channels, *Pestic. Biochem. Physiol.* 106 (2013) 101–112.
- [42] D.G. Heckel, Insecticide resistance after silent spring, *Science* 337 (2012) 1612–1614.
- [43] Z.D. Hu, X. Feng, Q.S. Ling, H.Y. Chen, Z.Y. Li, F. Yin, P. Liang, X.W. Gao, Biochemical mechanism of chlorantraniliprole resistance in the diamondback moth, *Plutella xylostella* Linnaeus, *J. Integr. Agr.* 13 (2014) 2452–2459.
- [44] J.G. Vontas, G.J. Small, J. Hemingway, Comparison of esterase gene amplification, gene expression and esterase activity in insecticide susceptible and resistant strains of the brown planthopper, *Nilaparvata lugens* (Stål), *Insect Mol. Biol.* 9 (2000) 655–660.
- [45] F.M. Chiang, C.N. Sun, Glutathione transferase isozymes of diamondback moth larvae and their role in the degradation of some organophosphorus insecticides, *Pestic. Biochem. Physiol.* 45 (1993) 7–14.
- [46] A.B. Dukre, M.P. Moharil, B.S. Ghodki, N.G.V. Rao, Role of glutathione S-transferase in imparting resistance to pyrethroids in *Plutella xylostella* (L.), *Int. J. Integr. Biol.* 6 (2009) 17–21.
- [47] C.H. Kao, C.N. Sun, In vitro degradation of some organophosphorus insecticides by susceptible and resistant diamondback moth, *Pestic. Biochem. Physiol.* 41 (1991) 132–141.
- [48] J. Hemingway, L. Field, J. Vontas, An overview of insecticide resistance, *Science* 298 (2002) 96–97.
- [49] S. Nehare, M.P. Moharil, B.S. Ghodki, G.K. Lande, K.D. Bisane, A.S. Thakare, U.P. Barkhade, Biochemical analysis and synergistic suppression of indoxacarb resistance in *Plutella xylostella* L., *J. Asia Pac. Entomol.* 13 (2010) 91–95.