



primary branch number (Li et al., 2016) in recent years. However, these potential loci identified with GWAS methods only explain a small amount of phenotypic variance (4.3~25.17% for seed quality traits, 16.47~20.51% for branch angle, 5.62~15.75% for flower time and 3.72~13.87% for yield-related traits). A probable reason for the low heritability detected with GWAS is that only additive models are applied but not considering dominance, epistasis (Zuk et al., 2012; Hemani et al., 2013). In a study of *Drosophila melanogaster* populations, about 50% of phenotypic variation in adult olfactory behavior was assigned to genotype-by-environment (G×E) interaction (Sambandan et al., 2008). Similarly, G×E or epistatic interactions could explain considerable proportion of variance of flowering traits in rice (Uwatoko et al., 2008) and in *Arabidopsis* (Caicedo et al., 2004; El-Lithy et al., 2006). Thus, trying to find the so-called “missing heritability” could help to efficiently dissect the genetic mechanism of complex traits (Manolio et al., 2009; Ingvarsson and Street, 2011; Resende et al., 2012).

In the earlier studies of QTL mapping, epistasis was observed for the resistance to *Sclerotinia sclerotiorum* in rapeseed, and the additive by additive interactions were the primary

0.140584(a)8.22229(b)0.e3335185(o)0.237963(e)-0.0625539(r)0.13994(a)8.0.24

i298(h)-G.83060

$$\begin{aligned}
& + \sum_i ae_{ih}u_{AE_{ih}} + \sum_i de_{ih}u_{DE_{ih}} + \sum_{i < j} aae_{ij}u_{AAE_{ij}} \\
& + \sum_{i < j} ade_{ij}u_{ADE_{ij}} + \sum_{i < j} dae_{ij}u_{DAE_{ij}} \\
& + \sum_{i < j} dde_{ij}u_{DDE_{ij}} + \varepsilon_h
\end{aligned} \quad (1)$$

where μ is the population mean; a_i is the additive effect of the i -th locus with coefficient u_{A_i} (1 for homozygote major alleles QQ and -1 for homozygote minor alleles qq); d_i is the dominance effect of the i -th locus with coefficient u_{D_i} (1 for heterozygote Qq 0 for homozygotes QQ and qq); aa_i , ad_i , da_i , and dd_i are the digenic epistasis effects with coefficients of random variables u_{AA_i} (1 for $QQ \times QQ$ and $qq \times qq$ -1 for $QQ \times qq$ and $qq \times QQ$), u_{AD_i} (1 for $QQ \times Qq$ -1 for $qq \times Qq$), u_{DA_i} (1 for $Qq \times QQ$ -1 for $Qq \times qq$), and u_{DD_i} (1 for $Qq \times Qq$); e_h is the random effect of the h -th environment; ae_{ih} is the additive \times environment interaction effect of the i -th locus in the h -th environment with coefficient $u_{AE_{ih}}$; de_{ih} is the dominance \times environment interaction effect of the i -th locus in the h -th environment with coefficient $u_{DE_{ih}}$; aae_{ij} , ade_{ij} , dae_{ij} , and dde_{ij} are the digenic epistasis \times race interaction effects in the h -th ethnic population with coefficients of random variables ($u_{AAE_{ij}}$, $u_{ADE_{ij}}$, $u_{DAE_{ij}}$, and $u_{DDE_{ij}}$); and ε_h is the residual effect of the i -th line or hybrid in the h -th environment.

Heritability of individual genetic effects were estimated by $h_g^2 = \alpha \sigma_g^2 / p$ ($\alpha = 2$ for additive effect, $\alpha = 1$ for dominant effect, $\alpha = 4$ for additive \times additive, $\alpha = 2$ for additive \times dominant or dominant \times additive, $\alpha = 1$ for dominant \times dominant), where phenotypic variance (p) is the sum of genetic variance (G), genetic by environment interaction variance (GE), and residual variance (ε),

$$\begin{aligned}
P &= G + GE + \varepsilon \\
&= (A + D + I) + (AE + DE + IE) + \varepsilon \\
&= (A + D + AA + AD + DA + DD) + \\
&\quad (AE + DE + AAE + ADE + DAE + DDE) + \varepsilon
\end{aligned} \quad (2)$$

The total heritability was estimated by

$$\begin{aligned}
h_{G+GE}^2 &= (h_A^2 + h_D^2 + h_I^2) + (h_{AE}^2 + h_{DE}^2 + h_{IE}^2) \\
&= (h_A^2 + h_D^2 + h_{AA}^2 + h_{AD}^2 + h_{DA}^2 + h_{DD}^2) \\
&\quad + (h_{AE}^2 + h_{DE}^2 + h_{AAE}^2 + h_{ADE}^2 + h_{DAE}^2 + h_{DDE}^2) \\
&= \sum_i h_a^2 + \sum_i h_d^2 + \sum_{i < j} h_{aa}^2 + \sum_{i < j} h_{ad}^2 + \sum_{i < j} h_{da}^2 \\
&\quad + \sum_{i < j} h_{dd}^2 + \sum_i h_{ae}^2 + \sum_i h_{de}^2 + \sum_{i < j} h_{aae}^2 + \sum_{i < j} h_{ade}^2 \\
&\quad + \sum_{i < j} h_{dae}^2 + \sum_{i < j} h_{dde}^2
\end{aligned}$$

We used the GMDR module (Generalized Multifactor Dimensionality Reduction; Qi et al., 2013) in the Q_XNetwor software (<http://ibi.zju.edu.cn/software/QTNetwork/>) to scan 33,689 SNP markers in 367 subjects for 1D~3D significant

candidate SNP markers, and obtained 539 candidate SNPs (260 in the A genome, 262 in the C genome, and 17 in the Scaffold group). The QTS mapping module in the Q_XNetwor was then used to dissect the genetic architecture of the eight agronomic traits of oilseed rape (Zhang et al., 2015). Significant SNPs associated with phenotypic variants were analyzed by setting a total of 2,000 permutation tests to calculate the critical P -value for controlling the experiment-wise type I error. The effects were predicted by using a Markov Chain Monte Carlo method with 20,000 Gibbs sampler iterations (Yang et al., 2007). The correlation coefficient ($R_{\hat{y}}$) between predicted breeding values and phenotypic values was estimated for each trait. Based on the predicted genetics effects of SNP loci for eight traits, we predicted total genotypic effects for the best lines and the best hybrids of the mapping population, and also predicted genotypic effects of superior lines and superior hybrids to inform further selection decisions (Yang and Zhu, 2005).

RESULTS

Estimated Heritability and Predicted Genetic Effects

We conducted a genome wide association study for eight yield-related traits of *B. napus* on a population including 151 inbred lines and 216 F₁ hybrids obtained from mating two female parents (L155 and L157) to other 149 inbred lines as male parents. Eight yield-related traits were investigated in our study by using a full genetic model with genetic effects of additive, dominance, epistasis, and their environment interactions. A total of 17 QTSs controlling eight yield traits were detected: 4 QTSs for PH, 2 QTSs for BN, 3 QTSs for IL, SS, ISN, TSW, BY, and SY, respectively (Figure 1 and Table S1). Some loci exhibited pleiotropic effects, including C09_M34850_G/A for six traits (BN, BY, ISN, PH, SS, and SY), A07_M11103_A/G for both BY and IL, A08_M12337_C/A for both PH and TSW, and C04_M26614_A/G for both BY and SY.

Estimated heritability and correlation coefficients ($R_{\hat{y}}$) between total genotypic values of detected QTSs and phenotypic values for the eight traits are listed in Table 1. The total heritability ranged from 58.47 to 87.98%, and was contributed by various types of genetic variance effects. With the exception of three yield traits (SS, BY, and SY), which were sensitive to the environment ($h_{GE}^2 \hat{=} 60.24\%$ for SY, 59.75% for BY, and 49.57% for SS), the other five traits were quite stable across the two environments ($h_{GE}^2 \hat{=} 4.14\sim 27.40\%$). With the exception of the SS trait, epistasis effects contributed a large portion of total heritability ($h_{I+IE}^2 \hat{=} 26.07\sim 62.14\%$). The correlation coefficient of genetic prediction with phenotype ($R_{\hat{y}} = 0.501\sim 0.899$) for each trait was very close to the estimated heritability ($h^2 \hat{=} 58.47\sim 87.98\%$), indicating that this statistic approach would be quite efficient for predicting the best lines/hybrids, and selecting the superior lines/hybrids by using the predicted genetic effects.

Highly significant (experiment-wise P_E -value $< 10^{-5}$) predicted genetic effects of 16 QTSs are presented in Table 1. Among the eight traits studied, plant height (PH) had the highest




FIGURE 1 | G x G plot of detected significant QTSs ($P_{EW} < 0.05$) for eight traits. Circle, QTS with additive effect; Square, QTS with dominant effect; Line between two QTSs, epistasis effect; Red color, QTS with general effects for two environments; Green color, QTS with environment-specific effects; Blue color, QTS with both general and environment-specific effects; Black color, QTS with significant epistasis effects but without detected individual effects.

TABLE 1 | Estimates of heritability and correlation coefficient of detected QTSs for eight traits.

Trait	h_A^2 (%)	h_D^2 (%)	h_I^2 (%)	h_{AE}^2 (%)	h_{DE}^2 (%)	h_{IE}^2 (%)	h_T^2 (%)	$R_{\hat{y}}$
PH	12.04	13.33	50.30	2.57	6.82	0.00	85.06	0.82
IL	4.15	20.75	53.01	3.92	0.00	6.15	87.98	0.73
BN	3.87	25.94	44.33	0.00	4.14	0.00	78.28	0.82
SS	0.00	8.90	0.00	35.91	13.66	0.00	58.47	0.50
ISN	3.86	35.52	5.79	3.03	0.00	24.37	72.57	0.74
TSW	17.23	8.32	15.11	2.15	7.57	10.96	61.34	0.62
BY	4.68	5.00	14.51	2.99	28.88	27.88	83.94	0.89
SY	1.27	11.11	13.96	5.51	6.55	48.18	86.58	0.90

h_A^2 , heritability of additive effects; h_D^2 , heritability of dominance effects; h_I^2 , heritability of epistasis effects (AA, AD, DA, and DD); h_{AE}^2 , heritability of environment-specific additive effects; h_{DE}^2 , heritability of environment-specific dominance effects; h_{IE}^2 , heritability of environment-specific epistasis effects (AAE, ADE, DAE, and DDE); h_T^2 , total heritability of all genetic effects. $R_{\hat{y}}$, correlation coefficient between phenotypic values and predicted genotypic values.

PH, plant height; IL, main inflorescence length; BN, branch number; SS, number of seeds per silique; ISN, effective silique number on main inflorescence; TSW, thousand seed weight; BY, biomass yield per plant; SY, seed yield per plant.

heritability ($h^2 \hat{=} 85.06\%$) mainly contributed by epistasis ($h_I^2 \hat{=} 50.30\%$). There was one locus (G/A of C09_M34850) detected with large and positive dominance effects. Homozygotes of this locus (G/G) could also increase plant height. The additive and dominance effects ($a \hat{=} -7.107$ of G/G, $d \hat{=} -3.565$ of G/A) were negative for locus Scaffold_M33906_G/A, which could be used in selecting for decreased plant height. Heterozygote C/A of A08_M12337 and homozygote G/G of C09_M34850 had a negative dominance \times additive epistasis effect ($da \hat{=} -11.003$), which could dramatically decrease plant height; however this large epistasis effect was counteracted by their main effects. Instead, the combination of homozygotes for major-allele C/C of A08_M12337 and minor-allele A/A of C09_M34850 implicated a lower plant height (Figure S2A). Epistasis effects were also the most important genetic effects on main inflorescence length (IL) ($h_I^2 \hat{=} 53.01\%$) and branch number (BN) ($h_I^2 \hat{=} 44.33\%$). Epistasis of heterozygote A/G of A03_M4640 \times minor-allele homozygote A/A of A08_M12338, and heterozygote A/G of A07_M11103 could significantly increase IL (Figures S2B,C). Heterozygote G/A of C09_M34850 was the major locus for increasing BN. And due to its epistasis effects, combination of heterozygote C/A of A01_M1364 \times minor-allele homozygote A/A of C09_M34850 could also significantly increase BN (Figure S2D).

Number of seeds per silique (SS) had very strong environment-specific additive and dominance effects ($h_{AE}^2 \hat{=} 35.91\%$ and $h_{DE}^2 \hat{=} 13.66\%$). Heterozygote G/A of A08_M12212 ($h_d^2 \hat{=} 6.43\%$), G/A of C09_M34850 ($h_d^2 \hat{=} 1.63\%$, $h_{de}^2 \hat{=} 5.95\%$) could increase SS in different environments.

Effective silique number on main inflorescence (ISN) had high heritability ($h^2 \hat{=} 72.57\%$) mostly due to dominance effects ($h_D^2 \hat{=} 35.52\%$) of three loci (A/G of A05_M8315, C/A of C04_M31883, and G/A of C09_M34850). Based on their large main effects, these three loci could be used in selection for increasing ISN, despite the negative dominance epistasis effect between heterozygotes of C04_M31883 and C09_M34850 (Figure S2E).

Thousand seed weight (TSW) had relatively large additive and epistasis variances ($h_A^2 \hat{=} 17.23\%$, $h_{I+IE}^2 \hat{=} 26.07\%$). Increasing TSW could be expected by homozygote minor-alleles A/A of A08_M12337 ($h_a^2 \hat{=} 15.03\%$), and also by heterozygote C/A of C01_M30818 ($h_d^2 \hat{=} 7.29\%$), but the heterozygote of both loci should be avoided due to negative epistasis effects ($dd \hat{=} -0.212$) (Figure S2F).

For biomass yield per plant (BY), the largest contributions of genetic variance were environment-specific dominance and

TABLE 2 | Predicted genetic effects of highly significant QTSs for eight traits.

Trait	Chr_SNP_Alleles	Effect	Predict	SE	-LogP _{EW}	h ² (%)
PH	A01_M114_G/A	d	2.63	0.56	5.70	0.48
	A08_M12337_C/A	a	-3.67	0.46	14.70	1.86
	C09_M34850_G/A	a	4.59	0.47	22.20	2.92
		d	13.09	0.43	201.20	11.86
		de1	6.80	0.60	28.70	6.21
	Scaffold_M33906_G/A	de2	5.21	0.60	17.30	6.21
		a	-7.11	0.46	52.20	6.99
	A08_M12337_C/A × C09_M34850_G/A	d	-3.57	0.43	16.00	0.88
da		-11.00	1.79	9.10	50.30	
IL	A07_M11103_A/G	d	5.42	0.25	105.70	15.01
		ae2	-1.94	0.37	6.70	2.26
	A08_M12338_G/A	a	-2.76	0.26	25.00	2.60
		d	3.35	0.25	41.50	5.74
	A10_M15289_A/C	a	1.64	0.27	9.20	0.91
	A03_M4640_A/G × A08_M12338_G/A	ad	-1.98	0.33	8.70	4.02
		da	-5.50	0.96	8.00	30.92
A07_M11103_A/G × A08_M12338_G/A	ad	-4.20	0.94	5.10	18.07	
BN	A01_M1364_C/A	a	-0.20	0.04	6.10	2.08
	C09_M34850_G/A	d	0.99	0.05	101.70	25.94
		de1	0.40	0.06	9.10	4.14
SS	A08_M12212_G/A	d	1.63	0.21	14.10	6.43
		ae1	1.09	0.18	9.00	6.90
	C01_M23080_G/A	ae1	-1.54	0.18	16.90	23.13
		d	0.82	0.15	7.60	1.63
		de2	1.47	0.21	12.00	5.95
ISN	A05_M8315_A/G	a	-1.71	0.34	6.40	2.57
		d	5.06	0.57	18.40	5.62
	C04_M31883_C/A	d	8.38	0.41	89.70	15.43
	C09_M34850_G/A	d	8.12	0.40	90.10	14.48
C04_M31883_C/A × C09_M34850_G/A		dde1	-4.54	0.60	13.20	9.04
TSW	A07_M11580_A/G	de2	0.18	0.03	8.70	4.81
	A08_M12337_C/A	a	-0.18	0.02	15.70	15.03
		de2	0.14	0.03	6.10	2.76
	C01_M30818_A/G	d	0.25	0.02	31.30	7.29
	A07_M11580_A/G × C01_M30818_A/G	dd	-0.21	0.02	18.30	10.74
BY	A07_M11103_A/G	a	-5.60	0.96	8.30	3.10
		d	8.96	0.92	21.60	1.98
		de2	23.75	1.32	71.50	27.84
	C04_M26614_A/G	d	7.79	0.95	15.60	1.50
	C09_M34850_G/A	d	7.85	0.91	17.10	1.52
SY	C04_M26614_A/G	d	2.10	0.28	13.60	2.16
	C09_M34850_G/A	d	4.26	0.27	56.90	8.94
		ae2	-2.37	0.40	8.60	5.51
		de2	4.74	0.38	35.20	5.96
	C04_M26614_A/G × C09_M34850_G/A	dde2	4.37	0.40	26.50	9.38

Chr_SNP_Alleles, genome chromosome_SNP_major allele/minor allele; *Effect*, genetic effect of QTS; *a*, additive effect for QQ locus; *-a* for qq locus; *d*, dominance effect for Qq locus; *dd*, dominance epistasis; *ae*₂, additive × environment interaction effect in Xiangyang; *de*₁, *de*₂, dominance × environment interaction effect in Wuhan, and Xiangyang, respectively; *dde*₂, dominance epistasis × environment interaction effect in Xiangyang; *Predict*, predicted genetic effect for the QTSs; *SE*, standard error of predicted effect; *-LogP_{EW}*, minus log₁₀(experiment-wise P-value); *h*², estimated heritability.

epistasis ($h_{DE}^2 \hat{=} 28.88\%$, $h_{IE}^2 \hat{=} 27.88\%$). Heterozygote A/G of A07_M11103 ($de_2 \hat{=} 23.745$, $h_{de_2}^2 \hat{=} 27.84\%$) could significantly increase BY (in environment Xiangyang). For the most important yield trait (seed yield per plant, SY), dominance and environment-specific epistasis ($h_D^2 \hat{=} 11.11\%$, $h_{IE}^2 \hat{=} 48.18\%$) were the major genetic recourses for increasing yield (Figure S2G). Heterozygote G/A of C09_M34850 could increase SY across environments ($h_d^2 \hat{=} 8.94$), and add extra selection response ($de_2 \hat{=} 4.743$) in environment Xiangyang ($h_{de_2}^2 \hat{=} 5.96\%$). Although dominance of heterozygote A/G of C04_M26614 could only slightly increase SY ($d \hat{=} 2.096$, $h_d^2 \hat{=} 2.16\%$), its epistasis interaction with dominance of another heterozygote G/A of locus C09_M34850 could also have large increase for SY in environment Xiangyang ($dde_2 \hat{=} 4.365$) (Figure S2H).

Predicted Genetic Effects for Different Genotypes

The Genotype-Phenotype (G-P) maps of epistasis SNPs for each trait, in both environments were presented in Figure S2. G-P maps exhibited various patterns as effects differed among different traits. High concordance was observed between

phenotypic G-P maps and genotypic G-P maps. Based on predicted genetic effects of QTSs for each trait, we further predicted the maximum and minimum genotypic effects of the superior lines and superior hybrids in two environments on eight traits. We also predicted the genotypic effects of homozygotes (QQ, qq), and heterozygote (Qq) for eight traits in two environments, respectively (Table 3). All the predicted genotypic effects of eight traits were negative for major-allele homozygote QQ, but positive for minor-allele homozygote qq. The predicted genotypic effects of heterozygote Qq were positive for all the eight traits studied. Among the eight traits for all-locus heterozygote (Qq) the predicted genotypic values were much larger than minor-allele homozygote (qq) for eight traits but not for TSW in one environment. It was implied that for this rapeseed population, hybrid breeding could potentially increase breeding values of seven yield traits but not for shortening plant height.

There was no difference between the best lines of mapping population and the predicted superior lines for four traits (BN, SS, TSW, and SY). It was suggested that pure-line variety breeding might have only limited potential for improving these traits based on the QTSs detected for this rapeseed population. For trait IL, breeding value of the best lines was smaller than the predicted superior lines, which was due to one locus in the best line L155 (minor-allele homozygote C/C of A01_M15289).

TABLE 3 | Predicted genetic effects in two environments for genotype of QQ, qq, Qq, best and superior lines, best and superior hybrids of eight traits.

Trait	Environment	Mean	QQ	qq	Qq	Best Line	Superior Line	Best Hybrid	Superior Hybrid
PH	G + GE1	133.7	-7.55	7.55	21.63	-7.55	-16.74	4.81	-16.74
	G + GE2	134.69	-5.06	5.06	15.6	-5.06	-19.22	2.32	-19.22
IL	G + GE1	52.02	-0.58	3.83	7.66	3.83	7.11	13.68	15.32
	G + GE2	50.85	-3.69	3.69	10.06	7.94	9.51	13.68	16.59
BN	G + GE1	4.77	-0.38	0.38	1.38	0.38	0.38	1.58	1.58
	G + GE2	7.92	-0.38	0.38	0.99	0.38	0.38	1.19	1.19
SS	G + GE1	23.44	-0.45	0.45	2.59	2.63	2.63	4.72	4.72
	G + GE2	22.66	-1.26	1.26	3.33	1.26	1.26	3.92	3.92
ISN	G + GE1	57.72	-1.69	1.69	15.27	9.64	4.15	15.27	15.27
	G + GE2	50.5	-5.24	5.24	19.81	7.88	5.24	19.81	19.81
TSW	G + GE1	3.69	-0.25	0.12	0.03	0.37	0.37	0.34	0.37
	G + GE2	3.38	-0.39	0.25	0.49	0.25	0.25	0.53	0.53
BY	G + GE1	51.47	-6.29	12.91	21.73	12.91	12.91	29.56	29.56
	G + GE2	92.9	-11.79	18.41	54.18	23.2	18.41	55.83	55.83
SY	G + GE1	10.79	-1.14	1.14	6.58	1.14	1.14	8.81	8.81
	G + GE2	23.79	-3.5	3.5	16.55	3.5	3.5	16.6	16.6

Mean, estimated mean of environment; E₁, 2013 in Wuhan; and E₂, 2013 in Xiangyang; QQ, homozygote of all loci with major-alleles; qq, homozygote of all loci with minor-alleles; Qq, heterozygote of all loci with Qq; Best line, predicted genotypic effect of line in the mapping population with lowest values for PH and highest values for other seven traits; Superior line, predicted genotypic effect of line in the selecting population with lowest values for PH and highest values for other seven traits; Best hybrid, predicted genotypic effect of hybrid in the mapping9658(h(e)-285#772869(pu)0.533#8(l)658(hl16(c)0.60133(tj)0.15#7(#n)0.0772869(c)0.601269(m)098237#(j)0.156#9(th)-280.227338)0.30#9(o)0.0772869(c)0.6017#(e)0.599#(j)0.126298(t)-2

The superior line could be obtained by replacing minor-alleles to major-alleles (A/A) of A01_M15289 for increasing $a \hat{=} 1.63$ in two environments and extra $ae_2 \hat{=} 1.27$ in E_2 . For another trait ISN, the breeding value of the best line ($G + GE \hat{=} 9.64$ in E_1 , and 7.88 in E_2) was larger than the superior line ($G + GE \hat{=} 4.15$ in E_1 , and 5.24 in E_2), because the best lines (L41 in E_1 and L45 in E_2) still had heterozygote locus (C/A of C04_M31883).


Among the eight traits studied, there were six traits (BN, SS, ISN, TSW, BY, and SY) with no difference of breeding values between the best hybrids and the superior hybrids in at least one environment. It was indicated that no selection advantage could be expected based on this mapping population for improving these six traits of hybrids. Expected gain of hybrid breeding could be obtained for two traits based on the best hybrids of this mapping population. For trait IL, the superior hybrid genotypes could be selected as A/G of A03_M4640 in two environments, and A/G of A07_M11103 in E_1 based on the best hybrids (L155 \times L76 in E_1 , L155 \times L26 in E_2) and maintained homozygote A/A for other two loci (A08_M12338, A01_M15289). There could have dramatic decrease for plant height of hybrid by just selecting all detected four QTSs of PH as major-allele homozygotes (C/C of A08_M12337, G/G of A01_M114, C09_M34850, and Scaffold_M33906).

The genotype of receptor lines and donor lines was listed in Table S2. Superior hybrid for six traits (BN, SS, ISN, TSW, BY, and SY) could be obtained via hybridization of acceptor lines (L155 and L157) to certain donor lines (L1~L154). There was only limited number of donor lines (3–13) that could contribute to target traits, except for trait ISN, which could be improved by 103 donor lines. We also found L102 was a competitive donor line of high potential. L102 could simultaneously improve four traits (SS, ISN, and SY) mainly due to its ability to donor G allele to A allele of C09_M34850, whose heterozygote could largely improve trait SS, ISN and SY with its pleiotropic effects. L102 also carried A/A of A07_M11580 and C/C of A08_M12337, which enabled L102 to improve TSW via introducing heterozygote into acceptor lines.

DISCUSSION

There are evidences that rare variants have large impacts on common human diseases (Cirulli and Goldstein, 2010; Yang et al., 2010; Zuk et al., 2014). Experimental evidence from association and linkage populations demonstrated that the rare genetic variation at β -carotene hydroxylase 1 (*crtRB*) was associated with β -carotene concentration in maize kernels (Yan et al., 2010). In our study, full genetic model including epistasis and environment-specific effects was firstly used to excavate the missing heritability and dissect genetic architecture of important agronomic traits in *B napus*. The average minor-allele frequency (MAF) in the 151 cultivars of this study was 11.07% (4.67~16.67%) for the detected 17 QTSs (10 A-alleles, 6 G-alleles, and 1 C-allele), and minor alleles had impacts on various genetic effects ($-a$ of qq , d of Qq , aa of $qq \times qq$, ad of $qq \times Qq$, $\pm da$ of $Qq \times qq$ and dd of $Qq \times Qq$) according to the genetic model. These minor alleles (qq) increased breeding values for all the eight traits and


made contributions to total heritability increasing by different genetic effects.

There were total 17 main additive effects loci and 14 environment-specific additive effects loci were identified, and most of them were negative (15 negative main additive effects and 8 negative environment-specific additive effects). However, the heritability of each additive locus was quite low ($h_a^2 \hat{=} 2.27\%$, $h_a^2 \hat{=} 0.12\sim 15.03\%$; $h_{ae}^2 \hat{=} 5.03\%$, $h_{ae}^2 \hat{=} 1.66\sim 23.13\%$). It suggested that the contribution to phenotype due to negative alleles could not be neglected and the additive variances were not the major genetic contribution for most traits studied. While previous study indicated that the additional effects of the alleles originated from both parents were detected to be important for yield components traits in rapeseed (Wang and Guan, 2010). The inconsistent of results reiterated the complex genetic mechanism of yield and yield-related traits, especially for allopolyploid species such as *B napus*. All the dominance variants were also contributed due to minor-allele heterozygotes (Qq). Dominance had much larger impacts on eight traits studied due to 21 main dominance loci with 19 having positive effects ($h_d^2 \hat{=} 6.14\%$, $h_d^2 \hat{=} 0.11\sim 25.94\%$) and 20 environment-specific dominance loci with 13 having positive effects ($h_{de}^2 \hat{=} 4.76\%$, $h_{de}^2 \hat{=} 0.30\sim 27.84\%$). Which indicated that positive dominance effects were conducive to yield-related traits plasticity in *B napus*. To data, epistatic effects were considered as important for complex traits in crops, such as plant height (Cao et al., 2001), yield (Huang et al., 2014), and salt tolerance (Wang et al., 2012) in rice seedlings, plant height in cultivated wheat (Zhang et al., 2008) and seed protein concentration in soybean (Qi et al., 2016). There were 10 pairs of loci identified with main dominance-related epistasis (dd , ad and da) ($h_i^2 \hat{=} 19.05\%$, $h_i^2 \hat{=} 1.35\sim 50.30\%$) and eight pairs of loci detected with environment-specific dominance-related epistasis (dde , ade and dae) ($h_{ie}^2 \hat{=} 14.11\%$, $h_{ie}^2 \hat{=} 0.74\sim 38.80\%$). The most contribution of heritability for yield traits was due to 12 pairs of loci identified with epistasis between dominance effects and additive effects (ad and da , ade and dae ; $h_{AD+DA+ADE+DAE}^2 \hat{=} 10.96\sim 53.01\%$). It was revealed the major role of epistasis influencing rapeseed yield. None of the detected QTSs was common with the SNPs detected in the previous association studies (Li F. et al., 2014; Li et al., 2016; Liu J. et al., 2016; Liu S. et al., 2016; Wang et al., 2016; Xu et al., 2016). It confirmed again that yield-related traits are complex polygenic phenomenon in rapeseed. Four pleiotropic QTSs were found to be associated with more than one trait (). It indicated that these traits might share part of genetic basis. Thus, the pleiotropic loci should be a priority for further research, and multi-traits should be taken into account together in genetic breeding practice.

Fully characterizing the genetic mechanism mediating heterosis is helpful for increasing crop yield. While none of the current genetic models can completely explain the heterosis phenomenon. Previous study indicated that epistasis together with all levels of dominance from partial to overdominance is responsible for the expression of heterosis in rapeseed (Radoev et al., 2008). In the study, large and positive heteroses for eight yield traits were mostly due to minor-alleles in heterozygotes

(Qq), and minor-allele homozygotes epistasis ($qq \times Qq$ of AD, $Qq \times qq$ of DA) was contributed for seven yield traits. It was concluded that epistasis together with heterozygotes loci play an important role in yield heterosis in *B. napus*.

Selection of best genotype combination is difficult due to the complexity of the genetic architecture. For example, to increase ISN, C/A of C04_M31883 \times G/A of C09_M34850 should be avoided due to their negative additive effects, but their large positive dominant effects could overturn the epistasis effects and make heterozygotes the optimal choice for these two loci. Genotype-Phenotype maps of epistasis SNPs based on prediction were adopted to visually demonstrate the accumulated genetic effects of the epistasis SNP pairs. Due to the high heritability of epistasis effects, the G-P maps based on population mean exhibited a similar pattern with corresponding G-P maps based on genetic prediction. But G-P maps based on population mean may be biased away from prediction due to confounding with effects of other loci and residual error. So G-P maps based on genetic prediction could be a better choice for selection by visualizing the true effects of epistasis effects. For loci involved in multiple epistases, selection needs more caution. For IL, A08_M12338 interacted with both A03_M4640 and A07_M11103. Although G/G of A07_M11103 combined with G/A of A08_M12338 could increase IL by $ad \hat{=} 4.204$, A/A of A08_M12338 should be chosen to obtain higher IL based on the total effects of the three loci. In this case, prediction function of superior line and superior hybrid is conducive to select the optimal genotype combination, thus efficiently utilize the heterosis in *B. napus*.

In the study, there were six traits (BN, SS, ISN, BY, TSW, and SY) having no difference of predicted breeding values between the best hybrids and the superior hybrids in at least one environment (). It was suggested that these six traits were already under strong breeding selection and conserved positive

effects of minor-allele homozygotes (qq) and heterexen f45(t)-09615057(e)1.

- Liu, J., Wang, W., Mei, D., Wang, H., Fu, L., Liu, D., et al. (2016). Characterizing Variation of Branch Angle and Genome-Wide Association Mapping in Rapeseed (*Brassica napus* L.). *Front Plant Sci* 7:21. doi: 10.3389/fpls.2016.00021
- Liu, R., Qian, W., and Meng, J. (2002). Association of RFLP markers and biomass heterosis in trigonometric hybrids of oilseed rape (*Brassica napus* x *B. campestris*). *heor App Genet* 105, 1050–1057. doi: 10.1007/S00122-002-1050-X
- Liu, S., Fan, C., Li, J., Cai, G., Yang, Q., Wu, J., et al. (2016). A genome-wide association study reveals novel elite allelic variations in seed oil content of *Brassica napus*. *heor App Genet* 129, 1203–1215. doi: 10.1007/s00122-016-2697-z
- Manolio, T. A., Collins, F. S., Cox, N. J., Goldstein, D. B., Hindorf, L. A., Hunter, D. J., et al. (2009). Finding the missing heritability of complex diseases. *Nature* 461, 747–753. doi: 10.1038/Nature08494
- Qi, H., Huang, J., Zheng, Q., Huang, Y., Shao, R., Zhu, L., et al. (2013). Identification of combining ability loci for five yield-related traits in maize using a set of testcrosses with introgression lines. *heor App Genet* 126, 369–377. doi: 10.1007/s00122-012-1985-5